

Carotid Endarterectomy

1. Endarterectomy for asymptomatic carotid artery stenosis. Executive Committee for the Asymptomatic Carotid Artery Stenosis Study. *JAMA*, 273:1421-1428, 1995. 2.
2. Badner NH, Beattie WS, Freeman D, Spence JD: Nitrous oxide-induced increased homocysteine concentrations are associated with increased postoperative myocardial ischemia in patients undergoing carotid endarterectomy. *Anesth Analg*. 91:1073-9, 2000.
3. Cheng MA, Theard MA, Tempelhoff R: Anesthesia for carotid endarterectomy: a survey. *J Neurosurg Anesthesiol* 9:211-216, 1997.
4. Fiori L, Parenti G: Electrophysiological monitoring for selective shunting during carotid endarterectomy. *J Neurosurg Anesthesiol* 7:168-173, 1995.
5. Fiori L, Parenti G, Marconi F: Combined transcranial doppler and electrophysiologic monitoring for carotid endarterectomy. *J Neurosurg Anesthesiol* 9:11-16, 1997.
6. Gross CE, Bednar MM, Lew SM, Florman JE, Kohut JJ: Preoperative volume expansion improves tolerance to carotid artery cross-clamping during endarterectomy. *Neurosurgery* 43:222-226; discussion 226-228, 1998.
7. Hamdan AD, Pomposelli FB Jr, Gibbons GW, Campbell DR, LoGerfo FW: Perioperative strokes after 1001 consecutive carotid endarterectomy procedures without an electroencephalogram: incidence, mechanism, and recovery. *Arch Surg* 134:412-415, 1999.
8. Haupt WF, Horsch S: Evoked potential monitoring in carotid surgery: a review of 994 cases. *Neurology* 42:835-838, 1992.
9. Heyer EJ, Adams DC, Moses C, Quest DO, Connolly ES: Erroneous conclusion from processed electroencephalogram with changing anesthetic depth. *Anesthesiology* 92:603-607, 2000.
11. Lam AM, Manninen PH, Ferguson GG, Nantau W: Monitoring electrophysiologic function during carotid endarterectomy: a comparison of somatosensory evoked potentials and conventional electroencephalogram. *Anesthesiology* 75:15-21, 1991.
12. McCarthy RJ, Nasr MK, McAteer P, Horrocks M: Physiological advantages of cerebral blood flow during carotid endarterectomy under local anaesthesia. A randomised clinical trial. *Eur J Vasc Endovasc Surg*. 24:215-21, 2002.
13. Messick JJ, Casement B, Sharbrough FW, Milde LN, Michenfelder JD, Sundt TJ: Correlation of regional cerebral blood flow (rCBF) with EEG changes during isoflurane anesthesia for carotid endarterectomy: critical rCBF. *Anesthesiology* 66:344-349, 1987.
14. Michenfelder JD, Sundt TM, Fode N, Sharbrough FW: Isoflurane when compared to enflurane and halothane decreases the frequency of cerebral ischemia during carotid endarterectomy. *Anesthesiology* 67:336-340, 1987.
15. North American Symptomatic Carotid Endarterectomy Trial Collaborators. Beneficial effect of carotid endarterectomy in symptomatic patients with high-grade carotid stenosis. *N Engl J Med* 325:445-453, 1991.

16. Pandit JJ, McLaren ID, Crider B: Efficacy and safety of the superficial cervical plexus block for carotid endarterectomy. *Br J Anaesth* 83:970-972, 1999.
17. Papavasiliou AK, Magnadottir HB, Gonda T, Franz D, Harbaugh RE: Clinical outcomes after carotid endarterectomy: comparison of the use of regional and general anesthetics. *J Neurosurg* 92:291-296, 2000.
18. Redekop G, Ferguson G: Correlation of contralateral stenosis and intraoperative electroencephalogram change with risk of stroke during carotid endarterectomy. *Neurosurgery* 30:191-194, 1992.
19. Sbarigia E, DarioVizza C, Antonini M, Speziale F, Maritti M, Fiorani B, Fedele F, Fiorani P: Locoregional versus general anesthesia in carotid surgery: is there an impact on perioperative myocardial ischemia? Results of a prospective monocentric randomized trial. *J Vasc Surg* 30:131-138, 1999.
20. Self DD, Bryson GL, Sullivan PJ: Risk factors for post-carotid endarterectomy hematoma formation. *Can J Anaesth* 46:635-640, 1999.
21. Stoneham MD, Doyle AR, Knighton JD, Dorje P, Stanley JC: Prospective, randomized comparison of deep or superficial cervical plexus block for carotid endarterectomy surgery. *Anesthesiology* 89:907-912, 1998.
22. Tangkanakul C, Counsell C, Warlow C: *Cochrane Database Syst Rev*; CD000126, 2000.
23. Wilhelm W, Schlaich N, Harrer J, Kleinschmidt S, Muller M, Larsen R: Recovery and neurological examination after remifentanil-desflurane or fentanyl-desflurane anaesthesia for carotid artery surgery. *Br J Anaesth.* 86:44-9, 2001.