

## **Transport Checklist**

- ☐ Airway
  - ☐ Extra intubation equipment
    - ☐ LMA(s) vs. I-Gel(s)
    - ☐ Macintosh 3/4 vs. Miller 2/3
    - ☐ ETT 6.0/7.0
    - ☐ Stylet(s)
    - ☐ 10cc syringe
  - ☐ Oral airway(s)
  - ☐ Tape
  - ☐ Consider bougie
  - ☐ Consider portable aspirator and suction catheter (Yankauer and bronchial)
- ☐ Breathing
  - ☐ Nonrebreather mask vs. nasal cannula
  - ☐ Ambu bag with PEEP valve
  - ☐ Jackson Reese, mask, bag
  - ☐ Transport ventilator for all intubated patients
  - ☐ CO2 detector (no CO2 monitoring on transport monitors nor ventilators)
  - ☐ Consider 2 full oxygen cylinders
- ☐ Circulation
  - ☐ Transport monitor detached from Perseus
  - ☐ Route for venous access isolated and secured
  - ☐ Consider IV kit with IV catheters
  - ☐ Adequate blankets to keep patient warm
- ☐ Medications
  - ☐ Sedation
    - ☐ Propofol
    - ☐ Other: \_\_\_\_\_
  - ☐ Analgesia
    - ☐ Fentanyl
    - ☐ Other: \_\_\_\_\_
  - ☐ Paralytics
    - ☐ Rocuronium
    - ☐ Succinylcholine
  - ☐ Vasoactive/antihypertensive
    - ☐ Phenylephrine
    - ☐ Ephedrine
    - ☐ Consider esmolol or labetalol
  - ☐ Emergency
    - ☐ 1 syringe box of epinephrine
    - ☐ 1 syringe box of atropine
    - ☐ Reversal agents
      - ☐ Sugammadex
      - ☐ Consider naloxone and flumazenil
  - ☐ Extra fluid-loading solution
  - ☐ Extra running infusions (vasoactive, antihypertensive, blood products)
  - ☐ Consider saline flushes
- ☐ Transport
  - ☐ Transport team: minimum of 3 escorts available including 1 anesthesia MD
  - ☐ Electrical reserve on transport equipment
  - ☐ Consider elevator key
  - ☐ Enter feet first into elevators during transport